



UNIVERSITI PUTRA MALAYSIA

**DEVELOPMENT OF OPTICAL BIOSENSOR TO DETERMINE
BENZOIC ACID CONCENTRATION BASED ON ITS INHIBITIVE
EFFECT ON TYROSINASE ENZYME**

HAIRUL HISHAM HAMZAH

FS 2010 29

**DEVELOPMENT OF OPTICAL BIOSENSOR TO DETERMINE BENZOIC ACID
CONCENTRATION BASED ON ITS INHIBITIVE EFFECT ON TYROSINASE
ENZYME**

HAIRUL HISHAM HAMZAH

**MASTER OF SCIENCE
UNIVERSITI PUTRA MALAYSIA**

2010



**DEVELOPMENT OF OPTICAL BIOSENSOR TO DETERMINE BENZOIC ACID
CONCENTRATION BASED ON ITS INHIBITIVE EFFECT ON TYROSINASE
ENZYME**

By

HAIRUL HISHAM HAMZAH

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, In
Fulfilment of the Requirements for the Degree of Master of Science**

September 2010



DEDICATIONS

To Assoc. Prof. Dr. Nor Azah Yusof,
for her patience, guidance, understanding and trust....

To Prof. Dato' Dr. Abu Bakar Salleh and Assoc. Prof. Dr. Fatimah Abu Bakar,
for their patience and trust....

To my beloved mom and family,
for their love, support and concern....

To all my friends,
for their support and understanding....

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the Degree of Master of Science.

**DEVELOPMENT OF OPTICAL BIOSENSOR TO DETERMINE BENZOIC ACID
CONCENTRATION BASED ON ITS INHIBITIVE EFFECT ON TYROSINASE
ENZYME**

By

HAIRUL HISHAM BIN HAMZAH

September 2010

Chairman : Associate Professor Nor Azah Yusof, PhD

Faculty : Science

An optical biosensor using phenol biosensor for the determination of benzoic acid based on inhibitive effect of benzoic acid on tyrosinase enzyme has been developed in this study. Two methods were developed which were based on in-solution tyrosinase and immobilized tyrosinase. Tyrosinase enzyme, phenol and 3-methyl-2-benzothiazolinone hydrazone (MBTH) were used as reagents for determination of benzoic acid. In-solution and immobilized tyrosinase as sensing element were successfully applied for the determination of benzoic acid.

In the detection of benzoic acid based on in-solution tyrosinase, the solution changed its color from dark maroon to light maroon depending on the concentration of benzoic acid, due to the inhibition of tyrosinase reaction by benzoic acid with response time of thirty minutes. The absorbance of products as the output of enzymatic reaction was detected by using uv-visible spectrophotometer with maximum absorbance at 504 nm. The biosensor demonstrated optimum activity at pH 7. The relative standard deviation (RSD) of the reproducibility of this method was

very good with RSD value of 1.91 %. The dynamic range of benzoic acid concentration was found to be between 50 ppm to 700 ppm with the detection limit (LOD) of 0.19 ppm. The kinetic parameters Michaelis-Menten constant (K_M) and maximum absorbance (A_{max}) in the absence and presence of benzoic acid showed the inhibition of benzoic acid on tyrosinase activity is competitive inhibitor with the inhibition constant (K_i) 90.9 ppm.

Test strip for detection of benzoic acid was developed by immobilizing tyrosinase, phenol and MBTH into filter paper using polystyrene as a polymeric support. The sensing scheme was based on the decreasing intensity of maroon color of test strip when dipped into benzoic acid solution. The test strip was characterized using optical fiber reflectance spectrometer (OFRS) and the result showed the test strip had a maximum reflectance at 375.65 nm. The optimum response of biosensor was achieved at pH 7. A linear response of the biosensor was obtained in the benzoic acid concentration range of 50 ppm to 700 ppm with LOD of 0.28 ppm. The reproducibility of the biosensor was good with calculated RSD of 0.47 %. The kinetic analyses showed that the inhibition of benzoic acid on tyrosinase activity was reversible and competitive inhibition with K_i of 52.9 ppm. The activity of the developed test strip was fairly sustained during for 20 days when stored at 3° C.

The results obtained from both methods developed in this study were compared with the established method of high performance liquid chromatography (HPLC). Excellent agreement was obtained between the developed and the HPLC methods.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Master Sains.

PEMBANGUNAN PENDERIABIO OPTIK UNTUK PENENTUAN KEPEKATAN ASID BENZOIK BERDASARKAN KESAN PERENCATANNYA KE ATAS ENZIM TIROSINASE

Oleh

HAIRUL HISHAM HAMZAH

September 2010

Pengerusi : Profesor Madya Nor Azah Yusof, PhD

Fakulti : Sains

Penderiablo optik menggunakan penderiablo fenol untuk penentuan asid benzoik berdasarkan kesan perencatan asid benzoik ke atas enzim tirosinase telah dibangunkan. Dua kaedah dibangunkan adalah yang mana berdasarkan tirosinase dalam larutan dan terpegun tirosinase. Enzim tirosinase, fenol, dan 3-metil-2-benzotiozolinon hidrazon (MBTH) digunakan sebagai bahan tindak balas untuk penentuan asid benzoik. Tirosinase dalam larutan dan terpegun sebagai elemen pengesanan telah berjaya diaplikasikan untuk penentuan asid benzoik.

Dalam penentuan asid benzoik berdasarkan tirosinase dalam larutan, larutan bertukar warna dari warna merah bata gelap kepada warna merah bata cerah bergantung kepada kepekatan asid benzoik merujuk kepada perencatan tindak balas tirosinase oleh asid benzoik dengan masa tindak balas selama 30 minit. Penyerapan produk sebagai hasil dari tindak balas tirosinase ditentukan menggunakan spektrofotometer ultra-nampak dengan penyerapan maksimum 504 nm. Penderiablo menunjukkan aktiviti optimum biosensor ialah pada pH 7. Sisihan piawai relatif

(RSD) bagi kebolehulangan kaedah ini adalah sangat baik dengan nilai RSD sebanyak 1.91%. Julat kepekatan dinamik bagi kepekatan asid benzoik ialah 50 ppm hingga 700 ppm dengan had pengesanan (LOD) adalah pada 0.19 ppm. Parameter kinetik untuk pemalar Michaelis-Menten (K_M) dan penyerapan maksimum (A_{max}) dalam ketidakhadiran dan kehadiran asid benzoik menunjukkan bahawa perencatan asid benzoik ke atas aktiviti tirosinase adalah perencat bersaing dan berbalik dengan pemalar perencatan (K_i) ialah 90.9 ppm.

Kepingan ujian untuk pengesanan asid benzoik dibangunkan dengan memegunkan tirosinase, fenol, dan MBTH ke dalam kertas turas menggunakan polistirina sebagai polimer sokongan. Skima pengesanan adalah berdasarkan pengurangan keamatan warna merah bata ke atas kepingan ujian apabila dicelup ke dalam larutan asid benzoik. Kepingan ujian ini dicirikan menggunakan spektrofotometer gentian optik (OFRS) dan keputusan menunjukkan bahawa kepingan ujian mempunyai pantulan maksimum pada 375.65 nm. Penderiablo mempunyai aktiviti optimum yang dicapai pada pH 7. Gerak balas linear penderiablo diperolehi dalam julat kepekatan asid benzoik adalah 50 hingga 700 ppm dengan had pengesanan adalah 0.28 ppm. Kebolehulangan penderiablo adalah baik dengan RSD ialah 0.47 %. Analisis kinetik menunjukkan bahawa perencatan asid benzoik ke atas enzim tirosinase adalah perencat bersaing dengan K_i ialah 52.9 ppm. Aktiviti kepingan ujian kekal selama 20 hari apabila disimpan pada suhu 3° C.

Keputusan yang diperolehi daripada kedua-dua kaedah yang dibina dalam kajian ini telah dibandingkan dengan kaedah yang diiktiraf iaitu kromatografi cecair prestasi tinggi (HPLC). Persetujuan yang sangat baik telah dicapai antara kaedah yang dibina dan kaedah HPLC.

ACKNOWLEDGEMENTS

In the name of Allah S.W.T, the most Merciful and the most gracious, who sends prophets for the guidance of mankind, million and billion of blessing upon the last prophet and seal of the prophets.

I would like to take this opportunity to express my sincere gratitude and appreciation to my supervisor, Associate Professor Dr Nor Azah Yusof for her guidance, assistance, encouragement, ideas and helpful critics throughout my work.

A very grateful thanks, I would like to express to my co-supervisors, Professor Dato' Dr. Abu Bakar Salleh and Associate Professor Dr. Fatimah Abu Bakar for their assistance, suggestion, unfailing help and constructive comments throughout the course of this project.

I would like to acknowledge the assistance and cooperation given by technical staff, Mr Isharudin Misron for the instrument of UV-visible spectrophotometer.

My appreciation also goes to Safura Taufik, Wan Siti Atikah, Shaari Daud, Kak Zue, and Lailatul Wahidah who have been helpful in sharing their knowledge, experience and cooperation and all my friends for their unforgettable support and concern toward the success of this project.

Finally, my special thanks are dedicated to my family for their never ending encouragement, support, advice, understanding and love.

I certify that a Thesis Examination Committee has met on 28 September 2010 to conduct the final examination of Hairul Hisham bin Hamzah on his Master of Science thesis entitled **“Development of Optical Biosensor to Determine Benzoic Acid Concentration Based on Its Inhibitive Effect on Tyrosinase Enzyme”** in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The committee recommends that the student be awarded the (Master of Science).

Members of the Thesis Examination Committee were as follows:

Name of Chairperson, PhD

Dr. Nor Azowa Ibrahim
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Name of Examiner 1, PhD

Associate Professor Dr. Irmawati Ramli
Faculty of Science
Universiti Putra Malaysia
(Internal Examiner)

Name of Examiner 2, PhD

Dr. Mohamed Ibrahim Mohamed Tahir
Faculty of Science
Universiti Putra Malaysia
(Internal Examiner)

Name of External Examiner, PhD

Professor Dr. Bahrudin Saad
School of Chemical Sciences
Universiti Sains Malaysia
(External Examiner)

BUJANG KIM HUAT, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:



This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Nor Azah Yusof, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Abu Bakar Salleh, PhD

Professor Dato'
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Member)

Fatimah Abu Bakar, PhD

Associate Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Member)

HASANAH MOHD GHAZALI, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

DECLARATION

I declare that the thesis is my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously, and is not concurrently, submitted for any other degree at Universiti Putra Malaysia or at any other institutions.

HAIRUL HISHAM HAMZAH

Date: 28 September 2010

TABLE OF CONTENTS

	Page
ABSTRACT	iii
ABSTRAK	v
ACKNOWLEDGEMENTS	vii
APPROVAL	viii
DECLARATION	x
LIST OF TABLES	xiv
LIST OF FIGURES	xv
LIST OF SCHEMES	xvii
LIST OF ABBREVIATIONS	xviii
 CHAPTER	
 I INTRODUCTION	 1
Biosensor	1
Principle of Enzyme-Based Biosensor	4
Optical Enzyme-Based Biosensor	6
Optical Transducer	7
Absorbance and Reflectance-Based Transducers	7
Enzyme Inhibition-Based Biosensor	10
Enzyme Immobilization	12
Technique and Support for Immobilization	13
Adsorption	14
Covalent Binding	15
Entrapment	17
Benzoic Acid	18
Toxicity Effect of Benzoic Acid on Human	20
Objective of This Research	21
 II LITERATURE REVIEW	 22
Review of Enzyme Inhibition-Based Biosensor	22
Biosensor-Based Test Strip	24
Analytical Methods for Determination of Benzoic Acid	29
Detection of Benzoic Acid Based on Inhibition Effect on Tyrosinase Enzyme	33
Benzoic Acid as Competitive Inhibitor	38



III	MATERIALS AND METHOD	41
	Chemical Reagents	41
	Instrumentations	42
	Ultraviolet-Visible Spectrophotometer	42
	Optical Fiber Spectrophotometer	42
	High Performance Liquid Chromatography (HPLC)	42
	Solution Preparations	43
	Preparation of Buffers	43
	Preparation of Reagents	43
	Preparation of Benzoic Acid Stock Solution	43
	Spectrophotometric Determination of Benzoic Acid Based on	
	In-Solution Tyrosinase	44
	In-Solution Tyrosinase	44
	Study on the Response Time	44
	Spectral Study	45
	pH Optimization	45
	Reproducibility of the Method	46
	Effect of Amount of Tyrosinase	46
	Effect of Different Concentration of Benzoic Acid	46
	Validation Study	47
	Interference Study	48
	Kinetic Study of Enzyme-Substrate Complex (ES)	48
	Kinetic Study of Enzyme-Inhibitor Complex (EI)	48
	Characterizations of Optical Test Strip Developed for Detection of	
	Benzoic Acid (Immobilized Tyrosinase)	49
	Preparation of Polystyrene Solution	49
	Preparation of Test Strip	49
	Characterizations of the Test Strip	50
	Study on the Response Time	50
	Spectral Study	50
	pH Optimization	50
	Reproducibility of the Test Strip	51
	Effect of Amount of Tyrosinase	51
	Effect of Different Concentration of Benzoic Acid	51
	Validation Study	52
	Interference Study	52
	Kinetic Study of Enzyme-Substrate Complex (ES)	52
	Kinetic Study of Enzyme-Inhibitor Complex (EI)	53
	Storage Stability of the Test Strip	53
IV	RESULTS AND DISCUSSION	54
	Spectrophotometric Determination of Benzoic Acid Based on	
	In-Solution Tyrosinase	54
	Study on the Response Time	54
	Spectral Study	54

	pH Optimization	55
	Reproducibility of the Method	57
	Effect of Amount of Tyrosinase	58
	Effect of Different Concentration of Benzoic Acid	59
	Validation Study	61
	Interference Study	62
	Kinetic Study	63
	Characterizations of Optical Test Strip Developed for Detection of Benzoic Acid (Immobilized Tyrosinase)	67
	Fabrication of the Test Strip	67
	Study on the Response Time	67
	Spectral Study	68
	pH Optimization	69
	Reproducibility of the Test Strip	70
	Effect of Amount of Tyrosinase	71
	Effect of Different Concentration of Benzoic Acid	72
	Validation Study	74
	Interference Study	75
	Kinetic Study	76
	Storage Stability of the Test Strip	78
V	CONCLUSIONS	80
	Recommendation for Further Studies	82
	REFERENCES	83
	APPENDICES	91
	BIODATA OF THE STUDENT	106